T S7/34

7/34/1
DIALOG(R)File 351:Derwent WPI
(c) 2005 Thomson Derwent. All rts. reserv.

009292971

WPI Acc No: 1992-420381/199251

Antimicrobial agent for plants esp. lawn grass - contains solvent extracts of plants, e.g. magnolia or salvia, natural additive e.g. chitosan and food additives e.g. alginic acid or sucrose fatty acid ester(s)

Patent Assignee: NAKANO SUMESE KK (NAKA-N)

Number of Countries: 001 Number of Patents: 002

Patent Family:

Patent No Kind Date Applicat No Kind Date Week JP 4316506 A 19921106 JP 91106370 Α 19910412 199251 B JP 3121036 B2 20001225 JP 91106370 19910412 200102 Α

Priority Applications (No Type Date): JP 91106370 A 19910412

Patent Details:

Patent No Kind Lan Pg Main IPC Filing Notes

JP 4316506 A 30 A01N-065/00

JP 3121036 B2 26 A01N-037/02 Previous Publ. patent JP 4316506

Abstract (Basic): JP 4316506 A

Antimicrobial agents are composed of one or more (a) one or more pressed juice, aq. and/or organic solvent extract or their condensates of Phellodendron amurense, Hosta undulata, Magnolia obovata, Aspidistra elatior, Scutellaria baicalensis, Rheum officinale, Symphytum officinale, Polygonum blumei, Celosia sp., Fragaria ananassa, bamboo, Rhus javanica, rose, nym, Thea sinensis, Castenea sp., Cinnamomum cassia, Stzygium aromaticum, Salvia officinalis, Peperomia sandersii and Humulus lupulus, (b) one or more natural additives of persimmon tannin, CaO, saponin, fumigating soln., naringin, hesperidin, leaf extract of Eucalyptus, benzoin resin, betaine, pectin degradation prod., soft roe protein, Monascus pigment, chitosan and polyysine, and (c) one or more food additives of lecithins, ferric lactate, tartaric acid, nicotinamide, alginic acid, fumaric acid, disodium 5'-guanylate, disodium orotate, sodium polyphosphate, sucrose fatty acid esters, thiamine laurylsulphate, thiamine thiocyanate, L-ascorbic acid, pyridoxine HCl, methylhesperidine, folic acid, sodium riboflavin 5'-phosphate, ferrous gluconate, ferric pyrophosphate, sodium benzoate, potassium sorbate, erythorbic acid, propylene glycol alginate, sodium metaphosphate, MgCl2, choline phosphate, glycine, L-alanine, adipic acid, vanillin, piperonal, sodium phytate, sorbitan monolaurate, tannic acid, gallic acid, propylene glycol, ethanol, lysozyme and sorbic acid, and a compsn. contg. acetic acid.

Pref. an acetic acid compsn. and one or more components (a), (b) and (c), pref. Magnolia obovata, Thea sinensis, Salvia officinalis, Syzygium aromaticum, Cinnamomum cassia, Humulus lupulus, ferric lactate and piperonal are mixed at ratios of 1:0.25-4 pts.wt. The resultant compsn. is diluted to give 0.002-0.2 wt.% of acetic acid and dispersed every five to 20 days.

USE/ADVANTAGE - Safe agricultural compsns. against harmful microorganisms, partic. fungi, without chemical injury.

In an example, a compsn. with various ratios of (a), (b), (c) and acetic acid was effective against fungi (e.g. Botrytis cenerea IFO 31831, Cladosporium carpophilum IFO 9645, Cl, fulvum IFO 8419, Fusarium oxysporum IFO 7152, Pythium aphanidermatum IFO 7030, Rhizoctonoia